Contact

Register

D



ISSN: 2157-7013

Journal of Cell Science & Therapy

Search.. search

3

in

Like us on: Like 39K

Make the best use of Scientific Research and information from our 700+ peer reviewed, Open Access Journals that operates with the help of 50,000+ Editorial Board Members and esteemed reviewers and 1000+ Scientific associations in Medical, Clinical, Pharmaceutical, Engineering, Technology and Management Fields.

Meet Inspiring Speakers and Experts at our 3000+ Global Conferenceseries Events with over 600+ Conferences, 1200+ Symposiums and 1200+ Workshops on

Medical, Pharma, Engineering, Science, Technology and Business

Review Article Open Access

Millimeter Wave and Drug Induced Modulation of the Immune System -Application in Cancer Immunotherapy

Mahendra K. Logani^{*}, Mahendra K. Bhopale and Marvin C. Ziskin

Center for Biomedical Physics, Temple University School of Medicine, USA

Conferences

*Corresponding Author: Dr. Mahendra K. Logani

Center for Biomedical Physics Temple University School of Medicine

3420 North Broad Street, Philadelphia, Pa 19140, USA

Tel: (215) 707-3321 Fax: (215) 707-4324

E-mail: mlogani@temple.edu

Received date: October 29, 2011; Accepted date: November 23, 2011; Published date: November 25, 2011

Citation: Logani MK, Bhopale MK, Ziskin MC (2011) Millimeter Wave and Drug Induced Modulation of the Immune System -Application in Cancer Immunotherapy. J Cell Sci Ther S5:002. doi: 10.4172/2157-7013.S5-002

Copyright: © 2011 Logani MK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Visit for more related articles at Journal of Cell Science & Therapy

View PDF Download PDF

Abstract

In recent years several approaches have been used for the treatment of cancer. These include surgery, chemotherapy, radiotherapy, and immunotherapy. Immunotherapy includes systemic administration of cytokines, adaptive transfer of activated T cells, NK cells, dendritic cells and macrophages. Immunotherapy is used to stimulate and train the patient's immune system to fight against cancer. However, this approach alone is not sufficient. Therefore, a combination of immunotherapy with conventional chemo- and radiation

Select your language of interest to view the total content in your interested language

Select Language 🗘

Powered by Google Translate

Post your comment

Share This Article











Relevant Topics

- Adaptive Cell Therapy
- Adaptive Immunity
- Adipose Derived Stem Cells (ASCs)
- Advanced Cell Therapy
- Advances in Anatomy
- Advances in Embryology
- Allergy Immunology
- Antibody-Producing Cells
- Antigen-Presenting Cells

Anontotic Dathwaye

Recommended Journals

therapy is commonly used for the treatment of cancer. But systemic administration of cytokines, such as interferons and interleukins, commonly used for immunotherapy, have their own toxic side effects that can be life threatening to many patients. In this article, we have reviewed the current knowledge of the effects of millimeter wave therapy (MMWT) on the immune system. MMWT, an alternative and complementary modality of treatment, is widely used for the treatment of cancer and many other diseases in Russia and several East European countries. However, it is virtually unknown to Western physicians. Our experimental studies have shown that MMWT when used in combination with chemotherapy protects the immune system from the toxicity of chemotherapy without introducing any additional toxicity of its own. Furthermore, our studies have shown that the combined millimeter wave- and chemotherapy can reduce the tumor metastasis, and tumor resistance to chemotherapeutic drugs. Thus combined therapy using MMWT provides a promising new strategy for the treatment of cancer.

Keywords

Millimeter wave therapy; T cells; NK cells; Macrophages; NF-κΒ; Metastasis; Chemoresistance; Apoptosis

Introduction

Millimeter electromagnetic waves (MMWs) are widely used for the treatment of many diseases in Russia, China, and many East European countries [1-6]. The three most common frequencies employed are 42.2, 53.6 and 61.2 GHz. These are approved by Russian Ministery of Health for medical treatment Excellent clinical results have been reported in the treatment of various diseases, including peptic ulcers, pain relief, cardiovascular diseases, wound healing, bronchial asthma, infantile cerebral palsy, skin disorders, chronic alcoholism, diabetic angioneuropathies, and cancer. Based upon some estimates, there are more than 1000 medical facilities in Russia alone where MMW therapy is being used, and the number of patients undergoing medical treatment exceeds 250 thousand per year [1]. In spite of the large numbers of patients treated with MMW therapy (MMWT) in the former Soviet Union, this treatment modality is virtually unknown to Western physicians and scientists. MMWs can be used as a monotherapy or in combination with other treatment methods. As an adjunct therapy, they are widely used to protect the immune system from the toxic effects of chemo- and radiation therapy in the treatment of cancer [3,7].

The usual treatment regimen consists of daily applications of low power MMW radiation for 15 to 30 minutes for 5 to 15 days. The MW device is typically a "book-sized" instrument which is brought in close contact with the skin surface. The site of application varies with the disease being treated. Surface wounds and skin diseases are usually treated at the site of the lesion. In treating arthritis, the site of application is at the affected joint. In treating internal diseases, the recommended site of application may be at any one of a number of anatomic or acupuncture points. A common site of application is the lower end of the sternum [6].

In recent years, several papers have been published on the biological effect of millimeter waves showing an increasing interest in the application, and understanding the mechanisms involved in MMWT [8-13]. In general, the effect of MMWT in clinical applications can be divided into three broad classes: (a) sedative and analgesic effects, (b) anti-inflammatory and repair stimulating action and (c) normalization of the immune system. There are many reports, mostly in the Russian literature, that MMWT produces both nonspecific and specific enhancement of the immune system [14,15]. The changes include an increase in phagocytic activity of macrophages [16], enhanced proliferation and normalization of the ratio of CD4+/CD8+ T-lymphocytes [17,18], and increased number of B-lymphocytes and normalized production of immunoglobulins [17-19]. It is generally believed that beneficial effects of MMWT in many diseases are due to enhancement of the immune response. In cancer treatment, MMWT has been used as monotherapy or in combination with chemo- and radiotherapy to increase immunity, and to reduce the toxic side effects of the above therapies [3,7,20]. Our experimental studies have shown that MMWT can reduce spread of melanoma metastasis [21]. In this short article, we review the current knowledge of the effects and molecular mechanisms of MMW

- Cell Signaling journal
- Cancer Research Journal
- Cell Science Journal
- Clinical and Laboratory
 Research Journal
- Clinical & Cellular Immunology Journal
- Molecular Biology Journal
- Cell Biology Journal
- Single Cell Biology Journal
- ▶ Stem Cell Research Journal

View More >>

Recommended Conferences

Cell Metabolism 2018
September 19 - 20, 2018
Philadelphia, USA

Article Tools

- Export citation
- Share/Blog this article

Article Usage

Total views: 12422

[From(publication date): specialissue-2012 - Sep 04, 2018]

Breakdown by view type

HTML page views: 8587

PDF downloads : 3835

therapy in protecting the immune functions from the toxic effects of chemotherapy. Specifically, we will discuss the effect of MMWT on T cells, NK cells and macrophage functions which constitute the most important cells in the fight against cancer.

Effect of MMWT on the T cell Functions

Our studies have shown that MMWT enhance T cell-mediated immunity. A delayed type hypersensitivity (DTH) assay in mouse skin was used to measure this effect [22]. Development of the DTH reaction in mouse or human skin has been shown to involve antigen-specific, T cell-mediated, memory dependent immunity [23,24]. Furthermore, we have examined whether MMWT can modulate T cell recovery after suppression with cyclophosphamide (CPA), an anticancer drug [25,26]. CPA is an alkylating agent that can prevent cell division by cross-linking DNA, leading to the reduction of tumor growth [27,28]. In addition to its cytotoxic effect on tumor cells, CPA is known to have multiple effects on the immune system resulting in a general depletion of immuno-competent cells [29]. A major side effect of this drug in chemotherapy is suppression of humoral and T cell-mediated immunity [30,31]. Among the problems associated with chemotherapy, restoration of immunocompetence is a critical issue to protect patients from secondary infections [30]. T cells have an impact on practically all aspects of immunity due to their ability to induce specific immune responses in both T and B cells. Also, T cells play an important role in the restoration process.

T cell functioning can be divided into an activation phase in which T cells divide and differentiate; and an effector phase, in which their function is expressed. Several basic effector functions have been defined for T cells: cytotoxic CD8⁺ T cells, directly kill infected cells, while, CD4⁺ -helper T (Th) cells can activate either B cells or macrophages in ways that are determined by the cytokines they produce. Proinflammatory cytokines, produced by Th1 cells, such as IFN-γ, activate, macrophages.and cytotoxic T cells; anti-inflammatory cytokines such as IL-10 produced by Th2 cells activate B cells. Our studies have shown that MMWs can restore production of IFN-γ and proliferation of T cells suppressed by CPA treatment. Conversely, no significant changes were observed in the level of IL-10 and proliferation of B cells. These results suggest that MMWs accelerate the recovery process selectively through a T cell-mediated immune response [25,26].

T cells play an important role in defense against tumor cells. Tumor cells can suppress functions of dendritic cells and macrophages thereby causing them to be incapable of stimulating and activating T cells sufficiently [32]. Considering the importance of these cells, clinical studies related to immunotherapy with activated T cells have been performed for various tumor types with promising results [33-35]. In view of our studies that MMWs can restore functions of T cells suppressed by chemotherapy, combined therapy should be very useful in cancer treatment.

Effect of Millimeter Wave Radiation on Natural Killer Cells

NK cells are known to kill a wide variety of tumor cells while sparing normal cells [36,37]. NK cells represent a distinct population of lymphocytes in terms of both phenotype and function [38]. They have large granular lymphocyte morphology and express characteristic cell surface receptors, such as the NK cell receptor protein-1 (termed NK1.1). NK cells mediate resistance to viral infections and cancer development and exhibit cytotoxic activity, which is nonrestricted to MHC (major histocompatibility complex) [39,40]. Thus, NK cells represent major effector cells of innate immunity. In addition, NK cells possess a variety of other functions, including the ability to produce a number of cytokines, involved in the modulation of hematopoiesis, immune responses and in the regulation of their own activities [41-43].

Our studies [44] have shown that treatment of mice with CPA, an anti-cancer drug, resulted in a significant suppression in cytolytic activity of NK cells. It was interesting to observe that irradiation of CPA treated mice with MMWs, accelerated the recovery of NK cell activity. It was further demonstrated that MMW irradiation significantly augment TNF- α production by NK cells suppressed by CPA administration. This observation supports the idea that the immunomodulatory effect of MMW irradiation could be beneficial when used in combination with chemotherapy. Similar results on the effect of MMWs on NK cell activity were reported by

Novoselova et al. [15] who studied the effect of MMWs on the immune system in mice with experimental tumors. MMW irradiation caused a significant increase in the production of TNF- α , nitric oxide and NK cell activity at the early stage of tumor development. MMW irradiation has also been shown to restore splenic NK cell activity suppressed by painful electric stimulation of the hind limbs in rats [45].

Several NK cell based immunotherapies have been developed in recent years [46], These include activation of a Patient's NK cells by injecting different cytokines such as IL-1,IL-2, IL-12 IL 15, and IL- 18 etc. Most commonly, IL-2 has been used for immunotherapy. It enhances NK cell-mediated IFN-γ production. The antitumor effect of IL-12 and IL-18 are also associated with upregulation of IFN-γ production by T cells and NK cells. IFN-γ has been shown to suppress tumor angiogenesis, and induce apoptosis in a variety of tumor cells by activating death receptors of the TNF super family. These include mainly FAS ligand receptors and TRAIL receptors [46]. However there are several limitations in the use of cytokines given exogenously for cancer treatment. Toxicity of systemic cytokine administration, and cytokine activated apoptosis of NK cells are two major limitations of cytokine mediated immunotherapies. Studies by us and others have shown that irradiation with millimeter waves increase the production of IFN-γ, TNF-α, IL-2 and enhance the cytotoxic activity of NK cells *in vivo* [44]. Therefore, immunotherapy by MMW should offer an advantage over the cytokine therapy given systemically. However, further experimental and clinical studies are needed to confirm this finding.

Effect of Millimeter Waves on Macrophage Functions

Macrophages are large mononuclear phagocytic cells important in innate as well as in cell mediated immunity. Macrophages play a critical role in host defense against microbial invasion and tumor cells. Macrophage activation is important in controlling of many key processes of immune system. When activated, macrophages are known to produce reactive oxygen and nitrogen species, free radicals, and TNF- α . TNF- α was first described as an endotoxin-induced and macrophage secreted factor that caused hemorrhagic necrosis of tumor cells. TNF- α is now recognized as a critical cytokine orchestrating cell differentiation, activation, proliferation and survival [47-50].

Our studies have demonstrated that CPA treatment of mice causes a strong decrease in spontaneous release of TNF- α by peritoneal macrophages that was restored to normal level when animals were irradiated with MMWs [26]. Several studies have shown that millimeter waves can activate macrophages [51]. The role of activated macrophages in the defense against tumor cells and in cancer treatment has been studied extensively [52]. It has been reported that macrophages can be involved in tumor cell killing, as well as in stimulation of tumor development and metastasis [53]. Activated macrophages are able to recognize, bind, and subsequently kill tumor cells. They can distinguish between tumor and normal cells based on differences in cell composition. Several studies have shown that a higher concentration in the amount of phosphotidylserine (PS) in the outer membrane of tumor cells is one of the factors responsible for specific tumor cell recognition and lysis or phagocytosis of tumor cells [52]. It has been shown that MMW irradiation can cause externalization of PS in the outer layer of membrane leading to apoptosis [54]. Since the rate of apoptosis depends upon the concentration of PS in outer layer of the cell membrane [55], MMWT holds a great potential to preferentially kill tumor cells because of greater accumulation of PS in their cell membrane.

Inhibition of Drug Resistance by MMWs

A major problem in cancer therapy is the development of tumor resistance to chemotherapeutic drugs. The major culprit involved in the development of drug resistance is the nuclear factor NF-κB, a transcription factor, that is involved in regulation of several genes. Many studies have shown that antineoplastic agents themselves can enhance this resistance through induction of NF-κB. There is ample evidence to indicate that both constitutive and inducible activation of NF-κB can protect tumor cells from apoptosis and thus enhance chemoresistance [56-58]. The above reports suggest that agents that inhibit activation of NF-κB by

chemotherapeutic drugs may also inhibit tumor resistance to chemotherapy. In support of the above hypothesis, it was shown that inhibition of inducible NF-kB activity reduces tumor resistance to chemotherapy [59,60].

Our studies have shown that MMW irradiation can inhibit activation of NF-κB induced by CPA, an anticancer drug [61]. These results suggest that MMW therapy when applied in conjunction with chemotherapy can reduce tumor resistance to the antineoplastic agents. The mechanisms through which MMWs inhibit drug induced activation of NF-κB remain to be elucidated.

Inhibition of Tumor Metastasis

One of the major side effects of chemotherapy is that although anticancer agents can reduce the growth of primary tumors, paradoxically, they may also enhance tumor metastasis. At the commonly used doses, most anticancer agents are immunosuppressive and therefore secondary neoplasias may result. For example, it has been reported that CPA can enhance tumor metastasis [62,63]. A recent study suggests that paclitexal, a drug curretly used for many forms of cancer might also increase metastasis [64]. It has also been shown that many antineoplastic agents including CPA can reduce NK cell activity [65,66]. Furthermore, it has been demonstrated that a reduction in NK cell activity results in enhancement of tumor metastasis [67-69]. Since our studies have demonstrated that the CPA induced suppression of NK cell activity can be restored by MMW therapy [44], we have further examined whether MMWs can also inhibit the enhanced tumor metastasis resulting from CPA pretreatment. In order to test this further, we used an 'experimental metastasis' model in which B16 melanoma cells were injected intravenously into C57BL/6 syngeneic mice and tumor development was evaluated in the lungs. The result of our studies showed that MMW irradiation at 42.2 GHz can reduce the metastasis enhanced by CPA treatment, suggesting that a combination of MMW-and chemotherapy can provide a promising new strategy to reduce metastasis in cancer treatment.


Molecular Mechanisms of MMW Treatment

We believe that biological effects of MMWs (penetration depth less than 1 mm in skin) are initiated by activation of free nerve endings in the skin [70-73]. Then the signal is conveyed to the central neural system where it modulates neural activity resulting in the development of various biological effects, such as the release of endogenous opioids. In support of this, it was shown that hypoalgesic effects of MMWs were abrogated when animals were treated with naloxone, a nonspecific blocker of opioid receptors, prior to MMW irradiation [74]. Furthermore, the involvement of endogenous opioids in suppression of melanoma growth by MMWs has also been reported [75].

In several studies from our group, it has been shown that MMW irradiation can induce release of endogenous opioids [76]. There is sufficient evidence that opioid peptides can modulate the immune system. The target of the effect of endogenous opioids could be NK cells [77], cytolytic functions of cytotoxic T cells, and the balance of Th1 and Th2 cytokines [78]. The disruption of a normal Th1/Th2 balance occurs in various immunological diseases [79,80]. In a recent study, we have reported that immunomodulation of T cells by millimeter waves is mediated by endogenous opioids [81]. However further studies are needed to determine whether endogenous opioids are involved in the observed effects of MMWs on the other components of the immune system suppressed by drug treatment. A systematic study on the combined effect of CPA and MMW on cytokine release in the presence or absence of naloxone and selective opioid receptor antagonists would be helpful in delineating the mechanisms involved in protection of immune functions by MMWs.

Conclusions and Perspectives

The studies reviewed above, suggest that MMW therapy may serve as an adjunct therapy to protect innate and adaptive immune functions from the adverse effects of chemotherapy in the treatment of cancer. Further in depth mechanistic studies in animal experiments and clinical trials, are needed to fully appreciate the value of millimeter wave therapy in combination therapy of human cancers. Specifically, more preclinical



studies need to be conducted to investigate the effect of MMWT on regulatory T cells, tumor cell growth, tumor apoptosis, tumor microenvironment, and tumor angiogenesis. In addition, the effect of MMWs on inflammatory enzymes which have been shown to enhance tumor growth and metastasis (e.g. cyclooxygenase 2, and metalloprotein-9) needs to be further investigated. The results of these studies will provide a scientific rationale for Western physicians to perform clinical trials in cancer patients.

Acknowledgements

Grant Support

The National Center for Complementary and Alternative Medicine/ NIH Grant numbers PO1-AT2002025 and RO1-AT00492.

References

- 1. Betskii OV, Devyatkov ND, Kislov VV (2000) <u>Low intensity millimeter waves in medicine and biology. Crit Rev Biomed Eng 28: 247-268.</u>
- 2. Rojavin MA, Ziskin MC (1998) Medical application of millimetre waves. QJM 91: 57-66.
- 3. Pletnev SD (2000) The use of millimeter band electromagnetic waves in clinical oncology. Crit Rev Biomed Eng 28: 573-587.
- 4. Pakhomov AG, Akyel Y, Pakhomova ON, Stuck BE, Murphy MR (1998) <u>Current state and implications of research on biological effects of millimeter waves: a review of the literature. Bioelectromagnetics 19: 393-413.</u>
- 5. Pakhomov AG, Mathur SP, Doyle J, Stuck BE, Kiel JL, et al. (2000) <u>Comparative effects of extremely high power microwave pulses and a brief CW irradiation on pacemaker function in isolated frog heart slices.</u>
 <u>Bioelectromagnetics 21: 245-254.</u>
- 6. Ziskin MC (2006) Physiological Mechanisms Underlying Millimeter Wave Therapy. In Current Concepts in Bioelectromagnetics Ayrapetyan S and Markov M (eds) Springer Pub 241-251.
- 7. Teppone M, Avakyan R (2010) Extremely high-frequency therapy in oncology. J Altern Complement Med 16: 1211-1216.
- 8. Wu G, Sferra T, Chen X, Chen Y, Wu M, et al. (2011) <u>Millimeter wave treatment inhibits the mitochondrion-dependent apoptosis pathway in chondrocytes. Mol Med Report 4:1001-1006.</u>
- 9. Beneduci A, Chidichimo G, De Rose R, Filippelli L, Straface SV, et al. (2005) <u>Frequency and irradiation time-dependant antiproliferative effect of low-power millimeter waves on RPMI 7932 human melanoma cell line.</u>
 <u>Anticancer Res 25: 1023-1028.</u>
- Beneduci A, Chidichimo G, Tripepi S, Perrotta E, Cufone F (2007) <u>Antiproliferative effect of millimeter radiation on human erythromyeloid leukemia cell line K562 in culture: ultrastructural- and metabolic-induced changes.</u>
 <u>Bioelectrochemistry 70: 214-220.</u>
- 11. Beneduci A (2009) Evaluation of the potential in vitro antiproliferative effects of millimeter waves at some therapeutic frequencies on RPMI 7932 human skin malignant melanoma cells. Cell Biochem Biophys 55: 25-32.
- 12. Li X, Du M, Liu X, Wu M, Ye H, et al. (2010) <u>Millimeter wave treatment inhibits NO-induced apoptosis of chondrocytes through the p38MAPK pathway. Int J Mol Med 25: 393-399.</u>
- 13. Li X, Du M, Liu X, Chen W, Wu M, et al. (2010) <u>Millimeter wave treatment promotes chondrocyte proliferation by upregulating the expression of cyclin-dependent kinase 2 and cyclin A. Int J Mol Med 26: 77-84.</u>
- 14. Safronova VG, Gabdoulkhakova AG, Santalov BF (2002) <u>Immunomodulating action of low intensity millimeter waves on primed neutrophils. Bioelectromagnetics 23: 599-606.</u>
- 15. Novoselova EG, Ogai VB, Sinotova OA, Glushkova OV, Sorokina OV, et al. (2002) Effect oh millimeter waves on the immune system in mice with experimental tumors. Biofizika 47: 933-942.
- 16. Rojavin M, Szabo I, Bussiere JL, Rogers TJ, Adler MW, et al. (1993) <u>Morphine treatment *in vitro* or *in vivo* decreases phagocytic functions of murine macrophages. Life Sci 53: 997-1006.</u>
- 17. Postovit NV, Shvets GL, Bondarenko SN, Kiichenko IuA (1989) Non-drug treatment of ulcers. Vrach Delo 2: 27-30.

- 18. Kutsenok VA (1994) The effect of electromagnetic radiation in the millimeter-wave range on the immune status of peptic ulcer patients. Lik Sprava 9-12: 139-142.
- 19. Babak OIa, Honcharova LI (1995) The microwave therapy of patients with duodenal peptic ulcer who took part in the cleanup of the aftermath of the accident at the Chernobyl Atomic Electric Power Station. Lik Sprava 7-8: 51-53.
- 20. Kabisov RK (2000) Millimeter radiation in the rehabilitation of oncological patients. Crit Rev Biomed Eng 28: 29-39.
- 21. Logani MK, Szabo I, Makar V, Bhanushali A, Alekseev S, et al. (2006) Effect of millimeter wave irradiation on tumor metastasis. Bioelectromagnetics 27: 258-264.
- 22. Logani MK, Yi L, Ziskin, MC (1999) <u>Millimeter waves enhance delayed-type hypersensitivity in mouse skin. Electro-and Magnetobiol 18: 165-176.</u>
- 23. Denkins Y, Fidler IJ, Kripke ML (1989) <u>Exposure of mice to UV-B radiation suppresses delayed hypersensitivity to Candida albicans</u>. Photochem Photobiol 49: 615-619.
- 24. Black CA (1999) Delayed type hypersensitivity: current theories with an historic perspective. Dermatol Online J 5: 7.
- 25. Makar V, Logani M, Szabo I, Ziskin M (2003) <u>Effect of millimeter waves on cyclophosphamide induced suppression of T cell functions. Bioelectromagnetics 24: 356-365.</u>
- 26. Makar VR, Logani MK, Bhanushali A, Alekseev SI, Ziskin M (2006) <u>Effect of cyclophosphamide and 61.22 GHz millimeter waves on T-cell, B-cell, and macrophage functions. Bioelectromagnetics 27: 458-466.</u>
- 27. Goodman M (2000) <u>Pentostatin (Nipent)</u> and <u>high-dose cyclophosphamide for the treatment of refractory autoimmune disorders. Semin Oncol 27: S67-S71.</u>
- 28. Allison AC (2000) <u>Immunosuppressive drugs: the first 50 years and a glance forward. Immunopharmacology 47:</u> 63-83.
- 29. Agarwal R, Diwanay S, Patki P, Patwardhan B (1999) <u>Studies on immunomodulatory activity of Withania somnifera</u> (<u>Ashwagandha</u>) <u>extracts in experimental immune inflammation</u>. <u>J Ethnopharmacol 67: 27-35.</u>
- 30. Angulo I, de las Heras FG, García-Bustos JF, Gargallo D, Muñoz-Fernández MA, et al. (2000) <u>Nitric oxide-producing CD11b(+)Ly-6G(Gr-1)(+)CD31(ER-MP12)(+)</u> cells in the spleen of cyclophosphamide-treated mice: implications for T-cell responses in immunosuppressed mice. Blood 95: 212-220.
- 31. Logani MK, Anga A, Szabo I, Agelan A, Irizarry AR, et al. (2002) <u>Effect of millimeter waves on cyclophosphamide induced suppression of the immune system. Bioelectromagnetics</u> 23: 614-621.
- 32. Rodrigues CM, Matias BF, Murta EF, Michelin MA (2011) <u>The role of T lymphocytes in cancer patients undergoing immunotherapy with autologous dendritic cells. Clin Med Insights Oncol 5: 107-115.</u>
- 33. Röllig C, Schmidt C, Bornhäuser M, Ehninger G, Schmitz M, et al. (2011) <u>Induction of cellular immune responses in patients with stage-I multiple myeloma after vaccination with autologous idiotype-pulsed dendritic cells.</u> <u>Immunother 34: 100-106.</u>
- 34. Santin AD, Bellone S, Palmieri M, Ravaggi A, Romani C, et al. (2006) <u>HPV16/18 E7-pulsed dendritic cell vaccination in cervical cancer patients with recurrent disease refractory to standard treatment modalities. Gynecol Oncol 100: 469-478.</u>
- 35. Matsuda K, Tsunoda T, Tanaka H, Umano Y, Tanimura H, et al. (2004) <u>Enhancement of cytotoxic T-lymphocyte</u> responses in patients with gastrointestinal malignancies following vaccination with CEA peptide-pulsed dendritic cells. Cancer <u>Immunol Immunother</u> 53: 609-616.
- 36. Smyth MJ, Hayakawa Y, Takeda K, Yagita H (2002) New aspects of natural-killer-cell surveillance and therapy of cancer. Nat Rev Cancer 2: 850-861.
- 37. Colucci F, Caligiuri MA, Di Santo JP (2003) What does it take to make a natural killer? Nat Rev Immunol 3: 413-425.
- 38. Trinchieri G (1989) Biology of natural killer cells. Adv Immunol 47: 187-376.
- 39. Vujanovic NL, Basse P, Herberman RB, Whiteside TL (1996) <u>Antitumor Functions of Natural Killer Cells and Control of Metastases</u>. <u>Methods 9: 394-408</u>.
- 40. Biron CA (1997) Activation and function of natural killer cell responses during viral infections. Curr Opin Immunol 9: 24-34.
- 41. Yu YY, Kumar V, Bennett M (1992) Murine natural killer cells and marrow graft rejection. Annu Rev Immunol 10:

189-213.

- 42. Djeu JY (1991) Natural killer cells. Role in resistance to cancer and infection. J Fla Med Assoc 78: 763-765.
- 43. Vitolo D, Vujanovic NL, Rabinowich H, Schlesinger M, Herberman RB, et al. (1993) Rapid Il-2-induced adherence of human natural killer cells. Expression of mRNA for cytokines and IL-2 receptors in adherent NK cells. J Immunol 151: 1926-1937.
- 44. Makar VR, Logani MK, Bhanushali A, Kataoka M, Ziskin MC (2005) <u>Effect of millimeter waves on natural killer cell activation</u>. <u>Bioelectromagnetics 26: 10-19.</u>
- 45. Shanin SN, Rybakina EG, Novikova NN, Kozinets IA, Rogers VJ, et al. (2005) <u>Natural killer cell cytotoxic activity and c-</u> Fos protein synthesis in rat hypothalamic cells after painful electric stimulation of the hind limbs and EHF <u>irradiation of the skin. Med Sci Monit 11: 309-315.</u>
- 46. Zamai L, Ponti C, Mirandola P, Gobbi G, Papa S, et al. (2007) NK cells and cancer. J Immunol 178: 4011-4016.
- 47. Gruss HJ, Dower SK (1995) <u>Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant</u> lymphomas. Blood 85: 3378-3404.
- 48. Locksley RM, Killeen N, Lenardo MJ (2001) <u>The TNF and TNF receptor superfamilies: integrating mammalian biology. Cell 104: 487-501.</u>
- 49. Chan FK, Chun HJ, Zheng L, Siegel RM, Bui KL, et al. (2000) A domain in TNF receptors that mediates ligand-independent receptor assembly and signaling. Science 288: 2351-2354.
- 50. Screaton G, Xu XN (2000) T cell life and death signalling via TNF-receptor family members. Curr Opin Immunol 12: 316-322.
- 51. Rojavin M, Szabo I, Bussiere JL, Rogers TJ, Adler MW, et al. (1993) <u>Morphine treatment *in vitro* or *in vivo* decreases phagocytic functions of murine macrophages. Life Sci 53: 997-1006.</u>
- 52. Klimp AH, de Vries EG, Scherphof GL, Daemen T (2002) A potential role of macrophage activation in the treatment of cancer. Crit Rev Oncol Hematol 44: 143-161.
- 53. Shih FF, Racz J, Allen PM (2006) <u>Differential MHC class II presentation of a pathogenic autoantigen during health</u> and disease. I Immunol 176: 3438-3448.
- 54. Szabo I, Kappelmayer J, Alekseev SI, Ziskin MC (2006) <u>Millimeter wave induced reversible externalization of phosphatidylserine molecules in cells exposed *in vitro*. Bioelectromagnetics 27: 233-244.</u>
- 55. Cummings M, Siitonen T, Higginbottom K, Newland AC, Allen PD (2002) <u>p53-mediated downregulation of Chk1</u> <u>abrogates the DNA damage-induced G2M checkpoint in K562 cells, resulting in increased apoptosis. Br J Haematol 116: 421-428.</u>
- 56. Beg AA, Baltimore D (1996) An essential role for NF-kappaB in preventing TNF-alpha-induced cell death. Science 274: 782-784.
- 57. Das KC, White CW (1997) <u>Activation of NF-kappaB by antineoplastic agents. Role of protein kinase C. J Biol Chem</u> 272: 14914-14920.
- 58. Hinz M, Löser P, Mathas S, Krappmann D, Dörken B, et al. (2001) <u>Constitutive NF-kappaB maintains high expression of a characteristic gene network, including CD40, CD86, and a set of antiapoptotic genes in Hodgkin/Reed-Sternberg cells. Blood 97: 2798-2807.</u>
- 59. Wang CY, Mayo MW, Baldwin AS Jr (1996) TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF-kappaB. Science 274: 784-787.
- 60. Uetsuka H, Haisa M, Kimura M, Gunduz M, Kaneda Y, et al. (2003) <u>Inhibition of inducible NF-kappaB activity</u> reduces chemoresistance to 5-fluorouracil in human stomach cancer cell line. Exp Cell Res 289: 27-35.
- 61. Logani MK, Natarajan M, Makar VR, Bhanushali A, Ziskin MC (2006) <u>Effect of millimeter waves on cyclophosphamide induced NF-kappaB. Electromagn Biol Med 25: 23-27.</u>
- 62. Carmel RJ, Brown JM (1977) <u>The effect of cyclophosphamide and other drugs on the incidence of pulmonary metastases in mice. Cancer Res 37: 145-151.</u>
- 63. Logani MK, Szabo I, Makar V, Bhanushali A, Alekseev S, et al. (2006) <u>Effect of millimeter wave irradiation on tumor metastasis</u>. <u>Bioelectromagnetics 27: 258-264</u>.

- 64. Quintavalle M, Elia L, Price JH, Heynen-Genel S, Courtneidge SA (2011) A cell-based high-content screening assay reveals activators and inhibitors of cancer cell invasion. Sci Signal 4: ra49.
- 65. Mantovani A, Luini W, Peri G, Vecchi A, Spreafico F (1978) Effect of chemotherapeutic agents on natural cell-mediated cytotoxicity in mice. I Natl Cancer Inst 61: 1255-1261.
- 66. Gazit Z, Kedar E (1994) <u>Chemotherapy-induced modulation of natural killer and lymphokine-activated killer cell activity in euthymic and athymic mice. Cancer Immunol Immunother 38: 243-252.</u>
- 67. Page GG, McDonald JS, Ben-Eliyahu S (1998) <u>Pre-operative versus postoperative administration of morphine:</u> impact on the neuroendocrine, behavioural, and metastatic-enhancing effects of surgery. Br J Anaesth 81: 216-223.
- 68. Page GG, Ben-Eliyahu S (2000) <u>Natural killer cell activity and resistance to tumor metastasis in prepubescent rats:</u>
 <u>deficient baselines, but invulnerability to stress and beta-adrenergic stimulation. Neuroimmunomodulation 7: 160-168.</u>
- 69. Page GG, Blakely WP, Ben-Eliyahu S (2001) <u>Evidence that postoperative pain is a mediator of the tumor-promoting effects of surgery in rats. Pain 90: 191-199.</u>
- 70. Alekseev SI, Ziskin MC, Kochetkova NV, Bolshakov MA (1997) <u>Millimeter waves thermally alter the firing rate of the Lymnaea pacemaker neuron. Bioelectromagnetics 18: 89-98.</u>
- 71. Pakhomov AG, Prol HK, Mathur SP, Akyel Y, Campbell CB (1997) <u>Search for frequency-specific effects of millimeter-wave radiation on isolated nerve function. Bioelectromagnetics 18: 324-334.</u>
- 72. Alekseev SI, Ziskin MC (1999) <u>Effects of millimeter waves on ionic currents of Lymnaea neurons.</u>
 <u>Bioelectromagnetics 20: 24-33.</u>
- 73. Alekseev SI, Ziskin MC (2003) <u>Local heating of human skin by millimeter waves: a kinetics study.</u>
 <u>Bioelectromagnetics 24: 571-581.</u>
- 74. Radzievsky AA, Gordiienko OV, Alekseev S, Szabo I, Cowan A, et al. (2008) <u>Electromagnetic millimeter wave induced hypoalgesia: frequency dependence and involvement of endogenous opioids. Bioelectromagnetics 29: 284-295.</u>
- 75. Radzievsky AA, Gordiienko OV, Szabo I, Alekseev SI, Ziskin MC (2004) <u>Millimeter wave-induced suppression of B16 F10 melanoma growth in mice: involvement of endogenous opioids. Bioelectromagnetics 25: 466-473.</u>
- 76. Rojavin MA, Cowan A, Radzievsky AA, Ziskin MC (1998) <u>Antipruritic effect of millimeter waves in mice: evidence for opioid involvement. Life Sci 63: PL251-257.</u>
- 77. Boyadjieva N, Dokur M, Advis JP, Meadows GG, Sarkar DK (2001) <u>Chronic ethanol inhibits NK cell cytolytic activity:</u> role of opioid peptide beta-endorphin. J Immunol 167: 5645-5652.
- 78. Sacerdote P, Gaspani L, Panerai AE (2000) The opioid antagonist naloxone induces a shift from type 2 to type 1 cytokine pattern in normal and skin-grafted mice. Ann N Y Acad Sci 917: 755-763.
- 79. Romagnani S (1994) Human TH1 and TH2 subsets: "eppur si muove"! Eur Cytokine Netw 5: 7-12.
- 80. Charlton B, Lafferty KJ (1995) The Th1/Th2 balance in autoimmunity. Curr Opin Immunol 7: 793-798.
- 81. Logani MK, Alekseev S, Bhopale MK, Slovinsky WS, Ziskin MC (2011) <u>Effect of millimeter waves and cyclophosphamide on cytokine regulation</u>. <u>Immunopharmacol Immunotoxicol 34: 107-112</u>.

Post your comment

me: *		
mail:		
ur comment: *		
ur comment: *		

Anti Spam Code:



C c

Can't read the image? click here to refresh

Post your comment

Peer Reviewed Journals

Make the best use of Scientific Research and information from our 700 + peer reviewed, Open Access Journals

Agri and Aquaculture	Geology & Earth Science
Biochemistry	Immunology & Microbiology
Bioinformatics & Systems Biology	Informatics
Biomedical Sciences	Materials Science
Business & Management	Mathematics
Chemical Engineering	Medical Sciences
Chemistry	Nanotechnology
Clinical Sciences	Neuroscience & Psychology
Computer Science	Nursing & Health Care
Economics & Accounting	Pharmaceutical Sciences
Engineering	<u>Physics</u>
Environmental Sciences	Plant Sciences
Food & Nutrition	Social & Political Sciences
General Science	Veterinary Sciences

Anesthesiology Molecular Biology Psychiatry Cardiology Nephrology Pulmonology Clinical Research Neurology Radiology Dentistry Nursing Reproductive Dermatology Nutrition Medicine Diabetes & Oncology Forderinglagy	Clinical & Medica	al Journals	
Gasteroenterology Genetics Pathology Haematology Pediatrics Healthcare Immunology Rehabilitation Infectious Diseases Medicine Microbiology	Cardiology Clinical Research Dentistry Dermatology Diabetes & Endocrinology Gasteroenterology Genetics Haematology Healthcare Immunology Infectious Diseases Medicine	Nephrology Neurology Nursing Nutrition Oncology Ophthalmology Orthopaedics Pathology Pediatrics Physicaltherapy &	Pulmonology Radiology Reproductive Medicine

International Conferences 2018-19

Meet Inspiring Speakers and Experts at our 3000+ Global Annual Meetings





Medical & Clinical Conferences		
Microbiology	Oncology & Cancer	
Diabetes & Endocrinology	Cardiology	
Nursing	<u>Dentistry</u>	
Healthcare Management	Physical Therapy Rehabilitation	
Neuroscience	<u>Psychiatry</u>	
Gastroenterology Genetics & Molecular Biology	Infectious Diseases Medical Ethics & Health Policies	
		<u>Pathology</u>
Alternative Healthcare	Reproductive Medicine & Women Healthcare	
<u>Pediatrics</u>	Surgery	
Ophthalmology	Radiology	

Conferences By Subject
Pharmaceutical Sciences
Pharma Marketing & Industry
Agri, Food & Aqua
Nutrition
Physics & Materials Science
Environmental Science
EEE & Engineering
<u>Veterinary</u>
Chemical Engineering
Business Management
<u>Massmedia</u>
Geology & Earth science

